

- [1] B.J. Dropesky, C.J. Orth, J. Inorg. Nucl. Chem. 24, 11 (1962)
- [2] T. Izak, S. Amiel, J. Inorg. Nucl. Chem. 34, 5 (1972)
- [3] L.-E. De Geer, G.B. Holm, Phys. Rev. C 22, 5 (1980)
- [4] H. Gausemel *et al.* Phys. Rev. C 69, 054307 (2004)
- [5] R. Lică *et al.* Phys. Rev. C 93, 044303 (2016)
- [6] P. Hoff *et al.*, Phys. Rev. Lett. 77, 6 (1996)
- [7] M. Piersa *et al.*, Phys. Rev. C 99, 024304 (2019)

Beta decay of ^{132}Cd : exploring the puzzle of decay half-lives southeast of ^{132}Sn

M Madurga¹, R Grzywacz^{1,2}, A Fijalkowska³, A Gottardo⁴, M Borge⁵, R Lica⁶

¹University of Tennessee, USA, ²Oak Ridge National Laboratory, USA, ³University of Warsaw, Poland, ⁴INFN, Italy, ⁵CSIC, Spain, ⁶National Institute for Physics and Nuclear Engineering, Romania

The region of nuclei around doubly magic ^{132}Sn [1] offers a unique window to study nuclear structure. Collective effects such as deformation quickly come into place a few proton numbers above tin [2]. On the other side of the $Z=50$ line, nuclei with proton number below Sn, such as Cd and Pd, are typically treated to be good shell model nuclei [3,4]. These calculations are in turn used to provide for nuclear parameters in models of isotope productions in stellar rapid neutron capture (r-process) [5]. However, recent measurements of the decay half-lives in the region show systematic discrepancies with shell model calculations. These discrepancies account for the majority of the deviations of calculated rprocess yields with solar abundances around the $A \sim 130$ peak [6]

In order to investigate the origin of the large discrepancies in beta-decay half-lives, the beta decay of $Z=48$, $N=84$ ^{132}Cd was studied at the ISOLDE facility, CERN. The large majority of the beta-decay strength was expected to populate the unbound lowest-energy 1^+ state through a Gamow-Teller transformation of a $g_{7/2}$ neutron into a $g_{9/2}$ proton [3,4], resulting in a (observed [7]) large neutron emission probability. In order to identify this state, the neutron time-of-flight array VANDLE [8,9] was installed at the ISOLDE decay station. The setup consisted in 26 VANDLE bars, for a 5% detection efficiency at 1 MeV, as well as 4 HPGe clovers, 3% efficiency at 1 MeV. We observed none of the gamma lines previously observed in ^{132}In in the decay of ^{133}Cd [10], nor any other gamma line that could be assigned to be in ^{132}In , suggesting a neutron branching ration close to 100%. Large Gamow-Teller strength was observed in the neutron time-of-flight spectra, as evidenced by 2/3 of the neutron intensity being emitted at 2 MeV. Analysis of the data indicates the Gamow-Teller strength occurs through 4 transitions to 4 individual 1^+ states in ^{132}In . The presence of multiple competing states, fragmenting the overall strength, offers a compelling explanation for the experimental half-life of ^{132}Cd being longer than state-of-the-art calculations [3]. The fragmentation of the 1^+ strength will be discussed in the framework of shell model and ab initio calculations using neutron-hole excitations of the ^{132}Sn core.

- [1] K.L. Jones et al., Nature 465, 454 (2010)
- [2] J.M. Allmond et al., Phys. Rev. Lett., 118, (2017) 092503
- [3] I.N. Borzov, Phys. Rev. C 71, 065801 (2005)
- [4] P. Möller, B. Pfeiffer, and K.-L. Kratz, Phys. Rev. C 67, 055802 (2003)
- [5] M.R. Mumpower, R. Surman, G.C. McLaughlin, and A. Aprahamian, Prog. Part. Nucl. Phys. 86, 86 (2016)
- [6] G. Lorusso et al., Phys. Rev. Lett. 114, 192501 (2015)
- [7] M. Hannawald et al., Phys. Rev. C 62, 054301 (2000)
- [8] W. A. Peters et al., Nucl. Inst. Meth. A836, 122 (2016)
- [9] S.V. Paulauskas et al., Nucl. Instrum Meth. A737, 22 (2014)
- [10] A. Jungclauss et al., Phys. Rev. C 93, 041301(R) (2016)